

Biologische Mediatoren zur Steuerung kieferorthopädischer Zahnbewegung

**Ein Beitrag von Dr. Ildeu Andrade Jr., Associate Professor in der Abteilung
Kieferorthopädie der Pontifícia Universidade Católica de Minas Gerais, Brasilien.**

Literatur

- 1) Krishnan V, Davidovitch Z. Cellular, molecular, and tissue-level reactions to orthodontic force. Am J Orthod Dentofacial Orthop 2006;129:469e.1-469e.32.
- 2) Pilon JJ, Kuijpers-Jagtman AM, Maltha JC. Magnitude of orthodontic forces and rate of bodily tooth movement. An experimental study. Am J Orthod Dentofacial Orthop. 1996 Jul;110(1):16-23.
- 3) van Leeuwen EJ, Maltha JC, Kuijpers-Jagtman AM. Tooth movement with light continuous and discontinuous forces in beagle dogs. Eur J Oral Sci. 1999 Dec;107(6):468-74.
- 4) Burstone CJ. The biomechanics of tooth movement. In: Kraus BS, Riedel RA, editors. Vistas in orthodontics. Philadelphia: Lea & Febiger; 1962.
- 5) Roberts WE, Huja S, Roberts JA. Bonemodeling—biomechanics, molecular mechanisms and clinical perspectives. Semin Orthod 2004;10:123-61.
- 6) Roberts WE, Ferguson DJ. Cell kinetics of periodontal ligament. In: Norton LA, Burstone CJ, editors. Biology of tooth movement. Boca Raton, Fla: CRC Press; 1995.
- 7) Rody WJ, King GJ, Gu G. Osteoclast recruitment to sites of compression in orthodontic tooth movement. Am J Orthod Dentofacial Orthop 2001;120:477-89.
- 8) Martin TJ, Ng K. Mechanisms by which cells of the osteoblast lineage control osteoclast formation and function. J Cell Biochem 1994;56:357-66.
- 9) Bozal CB, Fiol JA, Ubios AM. Early osteocyte response to bone resorption stimuli. Acta Odontol Latinoam 2001;14:24-9.
- 10) Drugarin D, Drugarin M, Negru S, Cioace R. RANK-RANKL/ OPG molecular complex—control factors in bone remodeling. TMJ 2003;53:296-302.
- 11) Kanzaki H, Chiba M, Takahashi I, Haruyama N, Nishimura M, Mitani H. Local OPG gene transfer to periodontal tissue inhibits orthodontic tooth movement. J Dent Res 2004;83:920-5.
- 12) Andrade I Jr, Taddei SRA, Souza PEA. Inflammation and tooth movement: the

- role of cytokines, chemokines and growth factors. *Semin in Orthod* 2012; 18:257-269.
- 13) Andrade Jr I, Silva TA, Silva GA, et al. The role of tumor necrosis factor receptor type 1 in orthodontic tooth movement. *J Dent Res* 2007;86:1089-1094.
- 14) Taddei SR, Andrade I Jr, Queiroz-Junior CM, Garlet TP, Garlet GP, Cunha Fde Q, Teixeira MM, da Silva TA. Role of CCR2 in orthodontic tooth movement. *Am J Orthod Dentofacial Orthop.* 2012 Feb;141(2):153-160.
- 15) Taddei SR, Queiroz-Junior CM, Moura AP, Andrade I Jr, Garlet GP, Proudfoot AE, Teixeira MM, da Silva TA. The effect of CCL3 and CCR1 in bone remodeling induced by mechanical loading during orthodontic tooth movement in mice. *Bone.* 2013 Jan;52(1):259-267.
- 16) Andrade Jr I, Taddei SRA, Garlet GP, et al. CCR5 down-regulates osteoclast function in orthodontic tooth movement. *J Dent Res* 2009;88:1037-1041.
- 17) Uematsu S, Mogi M, Deguchi T: Interleukin (IL)-1 beta, IL-6, tumor necrosis factor-alpha, epidermal growth factor, and beta 2-microglobulin levels are elevated in gingival crevicular fluid during human orthodontic tooth movement. *J Dent Res* 1996; 75:562-567.
- 18) Dai J, Rabie ABM: VEGF: An essencial mediator of both angiogenesis and endochondral ossification. *J Dent Res* 2007; 86:937-950.
- 19) Krishnan V, Davidovitch Z. On a path to unfolding the biological mechanisms of orthodontic tooth movement. *J Dent Res* 2009;88:597-608.
- 20) Teixeira CC, Khoo E, Tran J, Chartres I, Liu Y, Thant LM Khabensky I, Gart LP, Cisneros G, Alikhani M. Cytokine expression and accelerated tooth movement. *J Dent Res* 2010;89:1135-1141.
- 21) Fujita S, Yamaguchi M, Utsunomiya T, Yamamoto H, Kasa K. Low-energy laser stimulates tooth movement velocity via expression of RANK and RANKL. *Orthod Craniofac Res.* 2008 Aug;11(3):143-155.
- 22) Nishimura M, Chiba M, Ohashi T, Sato M, Shimizu Y, Igarashi K, Mitani H. Periodontal tissue activation by vibration: intermittent stimulation by resonance vibration accelerates experimental tooth movement in rats. *Am J Orthod Dentofacial Orthop.* 2008 Apr;133(4):572-583.
- 23) Kanzaki H, Chiba M, Arai K, Takahashi I, Haruyama N, Nishimura M, Mitani H. Local RANKL gene transfer to the periodontal tissue accelerates orthodontic tooth movement. *Gene Ther.* 2006 Apr;13(8):678-685.

- 24) Brooks PJ, Heckler AF, Wei K, Gong SG. M-CSF accelerates orthodontic tooth movement by targeting preosteoclasts in mice. *Angle Orthod.* 2011 Mar;81(2):277-283.
- 25) Dunn MD, Park CH, Kostenuik PJ, Kapila S, Giannobile WV. Local delivery of osteoprotegerin inhibits mechanically mediated bone modeling in orthodontic tooth movement. *Bone.* 2007 Sep;41(3):446-455.
- 26) Zhao N, Lin J, Kanzaki H, Ni J, Chen Z, Liang W, Liu Y. Local osteoprotegerin gene transfer inhibits relapse of orthodontic tooth movement. *Am J Orthod Dentofacial Orthop.* 2012 Jan;141(1):30-40.
- 27) Al-Qawasmi RA, Hartsfield JK Jr, Everett ET, Flury L, Liu L, Foroud TM, Macri JV, Roberts WE. Genetic predisposition to external apical root resorption. *Am J Orthod Dentofacial Orthop.* 2003 Mar;123(3):242-252.