

## **Implantation nach der Extraktion von Einzelzähnen im ästhetischen Bereich**

**Vivianne Chappuis, Daniel Buser**

<sup>1</sup> Hämmerle CH, Chen ST, Wilson TG Jr: Consensus statements and recommended clinical procedures regarding the placement of implants in extraction sockets. *Int J Oral Maxillofac Implants* 2004;19 Suppl: 26-28

<sup>1</sup> Chen ST, Buser D: Implant placement in post-extraction sites. Treatment options. Berlin: Quintessence Publishing, 2008

<sup>1</sup> Buser D, Maeglin B: Chirurgisches Vorgehen mit ITI-Implantaten. In *Orale Implantologie*, Schroeder A, Sutter F, Buser D, Krekeler G: Thieme: 260-328

<sup>1</sup> Schulte W, Kleineikenscheidt H, Lindner K, Schareyka R: The Tübingen immediate implant in clinical studies. *Dtsch Zahnärztl Z* 1978; 33: 348-359

<sup>1</sup> Chen ST, Wilson TG Jr, Hämmerle CH: Immediate or early placement of implants following tooth extraction: Review of biologic basis, clinical procedures, and outcomes. *Int J Oral Maxillofac Implants* 2004; 19 Suppl: 12-25

<sup>1</sup> Chen ST, Buser D: Clinical and esthetic outcomes of implants placed in post-extraction sites. *Int J Oral Maxillofac Implants* 2009; 24: (in press)

<sup>1</sup> Araújo MG, Lindhe J: Dimensional ridge alterations following tooth extraction. an experimental study in the dog. *J Clin Periodontol* 2005; 32: 212-8

<sup>1</sup> Araújo MG, Wennström JL, Lindhe J: Modelling of the buccal and lingual bone walls of fresh extraction sites following implant installation. *Clin Oral Implants Res* 2006; 17: 606-14

<sup>1</sup> Fickl S, Zuhr O, Wachtel H, Bolz W, Huerzeler MB: Hard tissue alterations after socket preservation: an experimental study in the beagle dog. *Clin Oral Implants Res* 2008; 19: 1111-8

<sup>1</sup> Aghaloo TL, Moy PK: Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement? *Int J Oral Maxillofac Implants* 2007; 22 Suppl: 49-70

<sup>1</sup> Buser D, Chen ST, Weber HP, Belser UC: Early implant placement following single-tooth extraction in the esthetic zone: biologic rationale and surgical procedures *Int J Periodontics Restorative Dent* 2008; 28: 441-51

<sup>1</sup> Jensen SS, Yeo A, Dard M, Hunziker E, Schenk R, Buser D: Evaluation of a novel biphasic calcium phosphate in standardized bone defects: a histologic and histomorphometric study in the mandibles of minipigs. *Clin Oral Implants Res* 2007 ;18 : 752-60

<sup>1</sup> Jensen SS, Brogini N, Hjørting-Hansen E, Schenk R, Buser D: Bone healing and graft resorption of autograft, anorganic bovine bone and beta-tricalcium phosphate. A histologic and histomorphometric study in the mandibles of minipigs. *Clin Oral Implants Res* 2006; 17:

237-43.

<sup>1</sup> Buser D, Weber HP, Lang NP: Tissue integration of non-submerged implants: 1-year results of a prospective study with 100 ITI hollow-cylinder and hollow-screw implants. Clin Oral Implants Res 1990; 1: 33-40

<sup>1</sup> Fürhauser R, Florescu D, Benesch T, Haas R, Mailath G, Watzek G: Evaluation of soft tissue around single-tooth implants crowns: the pink ethetic score. Clin Oral Implants Res 2005; 16: 639-33

<sup>1</sup> Belser UC, Grütter L, Vailati F, Bornstein MM, Weber HP, Buser D: Outcome evaluation of early placed maxillary anterior single-tooth implants using objective esthetic criteria: a cross-sectional, retrospective study in 45 patients with a 2- to 4-year follow-up using pink and whithe esthetic scores. J Periodontol 2009; 80: 140-51

<sup>1</sup> Choquet V, Hermans M, Adriaenssens P, Daelemans P, Tarnow DP, Malevez C: Clinical and radiographic evaluations of the papilla level adjacent to single-tooth dental implants. A retrospective study in the maxillary anterior region. J Periodontol 2001; 72: 1364-71

<sup>1</sup> Berglundh T, Lindhe J: Dimension of the periimplant mucosa. Biological width revisited. J Clin Periodontol. 1996; 23: 971-3

<sup>1</sup> Cochran DL, Hermann JS, Schenk RK, Higginbottom FL, Buser D: Biologic width around titanium implants. A histometric analysis of the implanto-gingival junction around unloaded and loaded nonsubmerged implants in the canine mandible. J Periodontol 1997; 68:186-98

<sup>1</sup> Kan JY, Rungcharassaeng K, Umezu K, Kois JC: Dimensions of the peri-implant mucosa: an evaluation of maxillary anterior single implants in humans. J Periodontol 2003; 74: 557-62

<sup>1</sup> Buser D, Martin W, Belser UC: Optimizing esthetics for implant restorations in the anterior maxilla: anatomic and surgical considerations Int J Oral Maxillofac Implants. 2004;19 Suppl:43-61

<sup>1</sup> Qahash M, Susin C, Polimeni G, Hall J, Wikesjö UM: Bone healing dynamics at buccal peri-implant sites. Clin Oral Implants Res 2008; 19: 166-72

<sup>1</sup> Cordaro L, Torsello F, Roccuzzo M: Clinical outcome of submerged versus non-submerged implants placed in fresh extraction sockets Clin Oral Implants Res 2009:(in press).

<sup>1</sup> Darby I, Chen S, De Poi R: Ridge preservation: what is it and when should it be considered. Aust Dent J 2008 ; 53: 11-21

Dental Tribune Swiss Edition Ausgabe September / 2009 Implant Special