

# White-Spot-Läsionen während Multibracketbehandlungen

Prof. Dr. Michael Knösel

DT German/Austrian/Swiss Edition 5/2014

---

Benson PE, Parkin N, Dyer F, Millett DT, Furness S, Germain P. Fluorides for the prevention of early tooth decay (demineralised white lesions) during fixed brace treatment. *Cochrane Database Syst Rev.* 2013 Dec 12;12:CD003809. doi:10.1002/14651858.CD003809.pub3.

Tufekci E, Dixon JS, Gunsolley JC, Lindauer SJ. Prevalence of white spot lesions during orthodontic treatment with fixed appliances. *Angle Orthod* 2011;81:206-10.

van der Veen MH, Attin R, Schwestka-Polly R, Wiechmann D. Caries outcomes after orthodontic treatment with fixed appliances: do lingual brackets make a difference? *Eur J Oral Sci.* 2010;118(3):298-303. doi: 10.1111/j.1600-0722.2010.00733.x.

Paris S, Hopfenmuller W, Meyer-Lueckel H. Resin infiltration of caries lesions: an efficacy randomized trial. *J Dent Res* 2010;89:823-6.

Knösel M, Bojes M, Jung K, Ziebolz D. Increased susceptibility for white spot lesions by surplus orthodontic etching exceeding bracket base area. *Am J Orthod Dentofacial Orthop* 2012;141:574-82.

Knösel M, Eckstein A, Helms H. Durability of esthetic improvement following Icon resin infiltration of multibracket-induced white spot lesions compared with no therapy over 6 months: a single-center, split-mouth, randomized clinical trial. *Am J Orthod Dentofacial Orthop* 2013;144 (1): 86-96.

Crombie F, Manton D, Palamara J, Reynolds E. Resin infiltration of developmentally hypomineralised enamel. *Int J Paediatr Dent.* 2013 Feb 15. doi: 10.1111/ipd.12025.

Munoz MA, Arana-Gordillo LA, Gomes GM, Gomes OM, Bombarda NH, Reis A, Loguercio AD. Alternative esthetic management of fluorosis and hypoplasia stains: blending effect obtained with resin infiltration techniques. *J Esthet Restor Dent.* 2013;25(1):32-9. doi: 10.1111/j.1708-8240.2012.00527.x.