

Literaturverzeichnis

4 Quadranten-Rehabilitation mit festsitzenden implantatgetragenen Brücken

Dr. Sven Egger, MSc, ZT Jürg Wermuth

- [1] Rufenacht CR
Fundamentals of Esthetics. Chicago:
Quintessenz, 1990: 67-134.

- [2] Jovanovic SA, Spiekermann H, Richter EJ.
Bone regeneration on titanium dental implants with dehisced defect sites. A clinical study.
Int J Oral Maxillofac Implants 1992 ; 7:233-245.

- [3] Bashutski JD, D'Silva NJ, Wang HL.
Implant compression necrosis: current understanding and case report.
J Periodontol 2009 Apr; 80(4):700-4.

- [4] Tarnow DP, Cho SC, Wallace SS.
The effect of interimplant distance on the height of inter-implant bone crest.
J Periodontol 2000; 71:546-549.

- [5] Adell R, Eriksson B, Lekholm U, Branemark P-I, Jemt T.
Long-term follow-up study of osseointegrated implants in the treatment of totally
edentulous jaws.
In J Oral Maxillofac Implants 1990; 5:347-359.

- [6] Monje A, Chan HL, Fu JH, Suarez F, Galindo-Moreno P, Wang HL.
Are Short Dental Implants (<10mm) Effective? A Meta-Analysis on Prospective Clinical
Trials.
J Periodontol. 2012; 23.

- [7] Esposito M, Cannizzaro G, Soardi E, Pistilli R, Piattelli M, Corvino V, Felice P.
Posterior atrophic jaws rehabilitated with prostheses supported by 6 mm-long, 4 mm-wide
implants or by longer implants in augmented bone. Preliminary results from a pilot
randomised controlled trial.
Eur J Oral Implantol. 2012 Spring; 5(1):19-33.