

Ausgabe: face 2/2014

Thema: Möglichkeiten und Grenzen der dermatologisch-ästhetischen Lasertherapie

Autoren: Dr. Peter Arne Gerber, Dr. Said Hilton

Literatur

1. Selective photothermolysis: precise microsurgery by selective absorption of pulsed radiation. **Anderson RR, Parrish JA.** Science. 1983 Apr 29;220(4596):524-7.
2. Comparison of the Q-switched alexandrite, Nd:YAG, and ruby lasers in treating blue-black tattoos. **Leuenberger ML, Mulas MW, Hata TR, Goldman MP, Fitzpatrick RE, Grevelink JM.** Dermatol Surg. 1999 Jan;25(1):10-4.
3. Lasertherapie der Haut. Raulin C, Karsai S, (2013). Springer, Berlin, Heidelberg
4. Pitfalls and recommendations in cases of laser removal of decorative tattoos with pigmented lesions: case report and **review** of the literature. Pohl L, Kaiser K, **Raulin C.** JAMA Dermatol. 2013 Sep;149(9):1087-9. doi: 10.1001/jamadermatol.2013.4901. **Review.**
5. Fractional photothermolysis: a new concept for cutaneous remodeling using microscopic patterns of thermal injury. Manstein D, Herron GS, Sink RK, Tanner H, Anderson RR. (2004). Fractional photothermolysis: a new concept for cutaneous remodeling using microscopic patterns of thermal injury. *Lasers Surg Med* **34**, 426-438.
6. Paasch U (2013). Fraktionale Laser: Wunsch und Wirklichkeit. *Akt Dermatol* **39**, 257-262.
7. Togsverd-Bo K, Haak CS, Thaysen-Petersen D *et al.* Intensified photodynamic therapy of actinic keratoses with fractional CO2 laser: a randomized clinical trial. *The British journal of dermatology* 2012; 166: 1262-9.