

Literatur

Chirurgische Behandlung der Eckzahndystopie

PD Dr. med. dent. habil. Dan Brüllmann

Oralchirurgie Journal 2/14

- 1) Paschos et al. Investigation of maxillary tooth sizes in patients with palatal canine displacement. *J Orofac Orthop.* 2005; 66:288-98.
- 2) Al-Nimri KS, Bsoul E. Maxillary palatal canine impaction displacement in subjects with congenitally missing maxillary lateral incisors. *Am J Orthod Dentofacial Orthop.* 2011 Jul;140(1):81-6.
- 3) Becker A. In defense of the guidance theory of palatal canine displacement. *Angle Orthod.* 1995;65(2):95-8.
- 4) Peck et al. The palatally displaced canine as a dental anomaly of genetic origin. *Angle Orthod.* 1994; 64: 249-256.
- 5) Sacerdoti R, Baccetti T. Dentoskeletal features associated with unilateral or bilateral palatal displacement of maxillary canines. *Angle Orthod.* 2004 Dec;74(6):725-32.
- 6) Al-Nimri KS, Bsoul E. Maxillary palatal canine impaction displacement in subjects with congenitally missing maxillary lateral incisors. *Am J Orthod Dentofacial Orthop.* 2011 Jul;140(1):81-6.
- 7) A. Becker, P. Smith, R. Behar The incidence of anomalous lateral incisors in relation to palatally displaced cuspids *Angle Orthod*, 51 (1981), pp. 24–29.
- 8) R. Sacerdoti, T. Baccetti. Dentoskeletal features associated with unilateral or bilateral palatal displacement of maxillary canines *Angle Orthod*, 74 (2004), pp. 725–732.
- 9) S. Peck, L. Peck, M. Kataja. The palatally displaced canine as a dental anomaly of genetic origin. *Angle Orthod*, 64 (1994), pp. 249–256.
- 10) S. Pirinen, S. Arte, S. Apajalahti. Palatal displacement of canine is genetic and related to congenital absence of teeth *J Dent Res*, 75 (1996), pp. 1742–1746.
- 12) Theodorakou C, Walker A, Horner K, Pauwels R, Bogaerts R, Jacobs R; SEDENTEXCT Project Consortium. Estimation of paediatric organ and effective doses from dental cone beam CT using anthropomorphic phantoms. *Br J Radiol.* 2012 Feb;85(1010):153-60. doi: 10.1259/bjr/19389412.
- 13) http://www.awmf.org/uploads/tx_szleitlinien/083-005_S1_Dentale_Volumentomographie_04-2009_12-2012.pdf
- 14) Boyd RL. Clinical assessment of injuries in orthodontic movement of impacted teeth. II. Surgical recommendations. *Am J Orthod.* 1984 Nov;86(5):407-18.
- 15) Michael E. Vermette, Vincent G. Kokich, and David B. Kennedy (1995) Uncovering labially impacted teeth: apically positioned flap and closed-eruption techniques. *The Angle Orthodontist*: February 1995, Vol. 65, No. 1, pp. 23-32.
- 16) Filippi et al. Sensory disorders after separation of the nasopalatine nerve during removal of palatal displaced canines: prospective investigation *British Journal of Oral and Maxillofacial Surgery*, 1999; 37: 134–136.
- 17) Chapokas et al. The impacted maxillary canine: a proposed classification for surgical exposure. *Oooo*. Vol. 113 No. 2 February 2012.
- 18) Joshi Transmigrant Mandibular Canines: A Record of 28 Cases and a Retrospective Review of the Literature. *The Angle Orthodontist* 2001; 71: 12-22.
- 19) Mupparapu Patterns of intraosseus transmigration and ectopic eruption of mandibular canines: review of literature and report of nine additional cases *Dentomaxillofacial Radiology* 2002; 31: 355-360.