

Ausgabe: face 2/2014

Thema: Wirkmechanismen und Anwendungsgebiete der Carboxytherapie

Autorin: Dr. Britta Knoll

Literatur

- O. Akça** et al., Hypercapnia improves tissue oxygenation. *Anaesthesiology* 2002, 97(4), 801-6
- O. Balik** et al., Does Carbon dioxide therapy really diminish localized adiposities? Experimental study with rats. *Aesthetic Plast Surg* 2011, 35(4), 470-4
- C. Brandi** et al., Carbon dioxide therapy in the treatment of localized adiposities. *Aesthetic Plast Surg* 2001, 25(3), 170-4
- C. Brandi** et al., Carbon dioxide therapy: effects on skin irregularity and its use as a complement to liposuction. *Aesthetic Plast Surg* 2004, 28(4), 222-5
- C. Brandi** et al., The role of Carbon dioxide therapy in the treatment of chronic wounds. *In Vivo* 2010, 24(2), 223-6
- C. Brandi** et al., Carbon dioxide may be not the only one but an efficient and secure gas for treating local adiposities. *Aesthetic Plast Surg* 2012, 36(1), 218-9
- M. Campana**, CO₂ therapy to treat adiposities and skin irregularities. *Prime-journal* 2013
- V. Campos** et al., Carboxytherapy for gynoid lipodystrophy treatment: The Brazilian Experience. *J Am Acad Dermatol* 2007
- JC. Ferreira** et al., Increase in collagen turnover induced by intradermal injection of Carbon dioxide in rats. *J Drugs Dermatol* 2008, 7(3), 201-6
- BR. Hartmann** et al., Effect of Carbondioxide enriched water and fresh water on the cutaneous microcirculation and oxygen tension in the skin of the foot. *Angiology* 1997, 48, 337
- GS Lee**, Carbon dioxide therapy in the treatment of cellulite: an audit of clinical practice. *Aesthetic Plast Surg* 2010, 34(2), 239-43
- T. Murohara** et al., Vascular endothelial growth factor/vascular permeability factor enhances vascular permeability via nitric oxide and prostacycline. *Circulation* 1998, 97(1),
- R. Nach** et al., Subcutaneous Carboxytherapy injection for aesthetic improvement of scars. *Ear Nose Throat J* 2010, 89(2), 64-6

Y. Sakai et al., A novel system for transcutaneous application of Carbon dioxide causing an artificial Bohr effect in the human body. PLoS ONE 2009, 6(9),e24137

V. Valaro et al., Carboxytherapy: effects on microcirculation and its use in the treatment of severe lymphedema. Acta Phleb 2007, 8(2), 79-91

U. Wollina et al., Transdermal CO₂ application in chronic wounds. Int J Low Extrem Wounds 2004, 3(2), 103-6

S. Zenker, Carboxytherapy - carbon dioxide injections in aesthetic medicine. Prime Journal, Jan/Feb. 2012, Vol 2, Issue 1