

Ausgabe: face 3/2014

Thema: Verbessertes Outcome in der Behandlung der Periorbitalregion – Filler und Botulinum richtig angewandt

Autorin: Dr. med. Marion Runnebaum

Literatur

1. Sadick NS, Bosniak SL, Cantisano-Zilkha M, Glavas IP, Roy D. Definition of the tear trough and the tear trough rating scale. *J Cosmet Dermatol.* 2007;6:218–22. [PubMed]
2. Haddock NT, Saadeh PB, Boutros S, Thorne CH. The tear trough and lid/cheek junction: Anatomy and implications for surgical correction. *Plast Reconstr Surg.* 2009;123:1332–40. [PubMed]
3. Belhaouari L, Teisseire P., Quinodoz P. Vectors for volumisation: enhancing the mid-face with HA. *PRIME* 2012;2(1):28-40
4. Allergan Zulassungsstudie, Fachinformation
5. Botulinum Toxin in Facial Rejuvenation: An Update. Jean Carruthers, MDa,*, Alastair Carruthers, MDb, *Dermatologic Clinics* 2009;27(4):417–425.
6. Meyer K. Palmer JW. The PS of the vitreous humor. *J Biol Chem.* 1934
7. Hirmand H. Anatomy and nonsurgical correction of the tear trough deformity. *Plast Reconstr Surg* 2010;125(2):699-708.8.Codner M
8. Carruthers et al, *Advances in Facial Rejuvenation: Botulinum Toxin Type A, Hyaluronic Acid Dermal Fillers, and Combination Therapies—Consensus Recommendations*, (*Plast. Reconstr. Surg.* 121 (Suppl.): 5S, 2008.)
9. Dormston et al, *Consensus Botulinum*, *JDDG*
10. Almeida AR et al. Glabellar wrinkles: A pilot study of contraction patterns. *Surg Cosmet Dermatol* 2010; 2(1): 23-8.