

Ausgabe: face 3/2014

Thema: Gesäßformung durch Bodylift-Operation

Autor: Dr. Harald Beck

Literatur

1. World Health Organization. Global Database on Body Mass Index. Available at: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html. Accessed 8/10/2014, 2014.
2. National Health and Nutrition Examination Survey 2009-2010
3. Post-Bariatric Buttock Contouring with Autogenous Tissue Augmentation
Zachary E. Gerut, Semin Plast Surg. 2006 February; 20(1): 30–37
4. Bilateral lumbar hip dermal fat rotation flaps: a novel technique for autologous augmentation gluteoplasty.
Raposo-Amaral CE, Cetrulo CL Jr, Guidi Mde C, Ferreira DM, Raposo-Amaral CM. Plast Reconstr Surg. 2006 May;117(6):1781-8.
5. Autologous gluteal augmentation after massive weight loss: aesthetic analysis and role of the superior gluteal artery perforator flap.
Colwell AS, Borud LJ.
Plast Reconstr Surg. 2007 Jan;119(1):345-56.
6. Autologous gluteal augmentation after massive weight loss.
Le Louarn C, Pascal JF.
Plast Reconstr Surg. 2008 Apr;121(4):1515-6; author reply 1516-7