

Ausgabe: face 4/2014

Thema: Volumisierung des Mittelgesichts

Autoren: Dr. Robert Kasten, Siegfried Schmidt

Literatur

- 1.** Alghoul M1, Codner MA. Retaining ligaments of the face: review of anatomy and clinical applications. *Aesthet Surg J.* 33:769-82. (2013)
- 2.** ASPS National Clearinghouse of Plastic Surgery Procedural StatisticsAmerican Society of Plastic Surgeons, Plastic Surgery Statistics Report (2012)
- 3.** Bernardin A et al. VYCROSS™: An innovative dermal filler technology; Posterpräsentation beim AMEC 2013.
- 4.** Carruthers J, Cohen SR, Joseph JH, Narins RS, Rubin M. The science and art of dermal fillers for soft-tissue augmentation. *J Drugs Dermatol.* 8:335-50. (2009)
- 5.** Eccleston D1, Murphy DK. Juvéderm(®) Volbella™ in the perioral area: a 12-month prospective, multicenter, open-label study. *Clin Cosmet Investig Dermatol.* 5:167-72 (2012)
- 6.** Funt D, Pavicic T. Dermal fillers in aesthetics: an overview of adverse events and treatment approaches. *Clin Cosmet Investig Dermatol.* 6:295-316. (2013)
- 7.** Gebrauchsanweisung zu Juvéderm® VOLBELLA® with Lidocaine 72526JR10 Stand 21.12.2012
- 8.** Gebrauchsanweisung zu Juvéderm® VOLIFT® with Lidocaine 72383JR11 Stand 24.09.2012.
- 9.** Gebrauchsanweisung zu Juvéderm® VOLUMA® with Lidocaine 72475JR11 Stand 19.09.2012.
- 10.** Ha RY, Nojima K, Adams WP Jr, Brown SA. Analysis of facial skin thickness: defining the relative thickness index. *Plast Reconstr Surg.* 115: 1769-73.(2005)
- 11.** Lambros V. Models of facial aging and implications for treatment. *Clin Plast Surg.* 35:319-27 (2008)
- 12.** Peter M. Prendergast. Anatomy of the Face and Neck, In: Cosmetic Surgery, M.A. Shiffmann and A. Di Guiseppe (Hrsg.) Springer Verlag, Berlin, Heidelberg S: 29-45 2012
- 13.** Philipp-Dormston WG1, Eccleston D, De Boulle K, Hilton S, van den Elzen H, Nathan M. A prospective, observational study of the volumizing effect of open-label aesthetic use of Juvéderm(®) VOLUMA(®) with Lidocaine in mid-face area. *J Cosmet Laser Ther.* 16:171-9. (2014)

- 14.** Rohrich RJ1, Arbique GM, Wong C, Brown S, Pessa JE. The anatomy of suborbicularis fat: implications for periorbital rejuvenation. *Plast Reconstr Surg.* 124:946-51. (2009)
- 15.** Sundaram H, Cassuto D. Biophysical characteristics of hyaluronic acid soft-tissue fillers and their relevance to aesthetic applications. *Plast Reconstr Surg.* 132:5S-21S. (2013)
- 16.** Taub AF, Sarnoff D, Gold M, Jacob C. Effect of multisyringe hyaluronic acid facial rejuvenation on perceived age. *Dermatol Surg.* 36:322-8 (2010)
- 17.** Wang F, Garza LA, Kang S, Varani J, Orringer JS, Gary J, Fisher GJ, Voorhees JJ. In Vivo Stimulation of De Novo Collagen Production Caused by Cross-linked Hyaluronic Acid Dermal Filler Injections in Photodamaged Human Skin. *Arch Dermatol.* 143:155-163 (2007)
- 18.** Wong CH1, Mendelson B. Facial soft-tissue spaces and retaining ligaments of the midcheek: defining the premaxillary space. *Plast Reconstr Surg.* 132: 49-56. (2013)
- 19.** Wulc AE, Sharma P, Czyz CN. The Anatomic Basis of Midfacial Aging In: M.E. Hartstein et al. (Hrsg.), *Midfacial Rejuvenation*, Springer Science+Business Media, LLC S. 15-28 (2012)