

## Literatur

### **Laserunterstützte Verbreiterung der keratinisierten Gingiva mit resorbierbarer 3D-Kollagenmatrix**

*Dr Michael Schäfer*

Laser Journal 3/14

- 1) Roos-Jansaker AM et al.: Nine-to fourteen-year follow-up of implant treatment. Part I: implant loss and associations to various factors. *J Clin Periodontol* 33, 283-289 (2006).
- 2) Roos-Jansaker AM et al.: Nine-to fourteen-year follow-up of implant treatment. Part II: presence of peri-implant lesions. *J Clin Periodontol* 33, 290-295 (2006).
- 3) Roos-Jansaker AM et al.: Nine-to fourteen-year follow-up of implant treatment. Part I: factors associated with peri-implant lesions. *J Clin Periodontol* 33, 296-301 (2006).
- 4) Block MS et al.: Hydroxyapatite-coated cylindrical implants in the posterior mandible: 10-year observations. *Int J Oral Maxillofac Implants* 11, 626-633 (1996).
- 5) Deporter DA et al.: A histological evaluation of a functional endosseous, porous-surfaced, titanium alloy dental implant system in the dog. *J Dent Res* 67, 1190-1195 (1988).
- 6) Kirsch A, Ackermann KL: The IMZ osteointegrated implant system. *Dent Clin North Am* 33, 733-791 (1989).
- 7) Araujo MG, Lindhe J: Dimensional ridge alterations following tooth extraction. An experimental study in the dog. *J Clin Peri- odontol* 2005;32: 212–218.
- 8) Araujo M, Linder E, Wennström J, Lindhe J: The Influence of Bio-Oss Collagen on Healing of an Extraction Socket: An Expe- rimantal Study in the Dog. *Int J Periodontics Restorative Dent* 2008;28:123–135.
- 9) Cardaropoli D, Cardaropoli G: Preservation of the postextracti- on alveolar ridge: a clinical and histologic study. *Int J Periodon- tics Restorative Dent.* 2008 Oct; 28(5):469–477.
- 10) Schropp L, Wenzel A, Kostopoulos L, Karring T: Bone healing soft tissue contour changes followig single-tooth extraction: A cli- nical and radiographic 12-month prospective study. *Int J Peri- odontics Restorative Dent* 2003;23:313–323.
- 11) Strietzel FP, Shakibaie-M B: Der Einsatz des TefGen-FD®-Mem- bran zum Erhalt des Alveolarkamms nach Zahnextraktionen: eine klinische Studie. *Dtsch Zahnärztl Z* 1998;12:883–886.
- 12) McGuire, M.K. & Scheyer, E.T. (2010): Xenogeneic collagen matrix with coronally advanced flap compared to connective tissue with coronally advanced flap for the treatment of dehiscence- type recession defects. *Journal of Periodontology* 81: 1108–1117.
- 13) Sanz, M., Lorenzo, R., Aranda, J.J., Martin, C. & Or- sini, M. (2009): Clinical evaluation of a new colla- gen matrix (Mucograft prototype) to enhance the width of keratinized tissue in patients with fixed prosthetic restorations: a randomized prospective clinical trial. *Journal of Clinical Periodontology* 36: 868–876.