

Literatur

Änderung der skelettalen Strukturen zur Harmonisierung der Weichteilstrukturen im Gesicht

Prof. Dr. med. Dent. Nezar Watted*, Dr. Muhamad Abu-Hussein**, Ali Watted***, Prov.Doz. Dr. Dr. Josip Bill****, Prof. Dr. Dr. med. Peter Proff***** , Dr. Dr. Benjamin Schlomi*****

Oralchirurgie Journal 1/2015

* Klinik und Polikliniken für Zahn-, Mund- und Kieferkrankheiten der Julius-Maximilians-Universität Würzburg, Germany

** Department of Pediatric Dentistry, University of Athens, Greece

*** Dental school of the university of Regensburg, Germany

**** Clinic for Cranio-maxillo-facial surgery Würzburg, Germany

***** Department of Orthodontics, University of Regensburg, Germany

***** Department of Oral and Maxillofacial Surgery, Saurasky Medical Centar and Tel Aviv University, Israel

1. Akay, M.C., Aras A. & Günbay T., et al.: Enhanced Effect of Combined Treatment with Corticotomy and Skeletal Anchorage in Open-bite Correction. *J Oral Maxillofac Surg.*, 2010, 67: 563-569
2. Aziz, S.R.. Simon P. Hullinen and the Origin of Orthognathic Surgery. *Journal of Oral and Maxillofacial Surgery*, 2004, 62: 1303-1307
3. Babcock W.W.: Surgical treatment of certain deformities of jaw associated with malocclusion of teeth. *J. Amer. Med. Ass.* 53: 833-839, 1909
4. Baek, S.H.; Ahn, H.W.; Kwon, Y.H. & Choi, J.Y.: Surgery First Approach in Skeletal Class III Malocclusion Treated with 2-Jaw Surgery: Evaluation of Surgical Movement and Postoperative Orthodontic Treatment. *Journal of Craniofacial Surgery*, 2010, .21: 332-338
5. Bell WH (1969) Revascularziation and bone healing after anterior maxillary osteotomy: A study using adult rhesus monkeys. *J Oral Surg* 27:249
6. Bell W.H.: Le Fort I Osteotomy for correction of maxillary dformities. *J. Oral Surg.* 33: 412-426, 1975
7. Bell WH, Proffit HR, White RP. Surgical correction of facial deformities. Saunder Verlag, Baltimor, 1980.
8. Bell WH, Schdeman GB. Correction of vertical maxillary deficiency: Stability and soft tissue changes. *J Oral surg* 1981; 39: 666-670.
9. Bell HW, Jacobs JD. Simultaneous repositioning of the maxilla, mandible, and chin. Treatment planning and analysis of the soft tissues. *Am J Orthod.* 91: 28-50, 1986.
10. Buddgaard M, Melsen B, Terp S. Changes during and following total maxillary osteotomy (Le Fort I Procedure): a cephalometric study. *Eur J Orthod* 1986; 8: 21-29.
11. Burstone GJ. The integumental profile. *Am J Orthod* 1958; 44: 1-25.

12. Burstone GJ. Lip posture and its significance in treatment planning. Am J Orthod 1967; 53: 262-84.
13. Calangna L, Silverman S, Garfinkel L. Influence of neuromuscular conditioning on centric relation registrations. J Prosthet Dent 1973; 30: 598-604.
14. Cheever DW: Nasopharyngeal Polypus, attached to the Basilar Process of Occipital and Body of the Sphenoid Bone successfully removed by a Section, Displacement, and subsequent Replacement and Reunion of the Superior Maxillary Bone. Boston Med Surg J 8:162-171, 1867
15. Celenza, F V. The centric position. replacement and character. J Prosthet Dent 1973; 30: 591-598.
16. Carlson DS, Ellis E, Dechow PC. Adptation of the suprathyroid muscle complex to mandibular advancement surgery. Am J Orthod 1987; 92: 134-143.
17. Coh-Stock C. Die chirurgische immediate Regulierung der Kiefer, chirurgische Behandlung der Prognathie. Dtsch Vjschr Zahnheik 1921; 37: 320-331.
18. Collins P, Epker BN. The alar basse Cinch: A technique for prevention of alar base flaring secondary to maxillary surgery. Oral Surg 1982; 53: 549-554.
19. Cunningham G : Methode sofortiger Regulierung von anomalen Zahnstellungen. Österr Vjschr Zahnheilk 10: 455-457, 1984
20. Dal Pont G. L`osteotomia retromolare per la correzione della progenia, Minerva chir., 1959; 18: 1138-1141.
21. Dal Pont G. Die retromolare Osteotomie zur Korrektur der Progenie, der Retrogenie und des Mordex apertus. Öst Z Stoma 1961; 58: 8-10.
22. Dann JJ, Fonseca RJ, Bell WH. Soft tissue changes associated with total maxillary advancement: a preliminary study. J Oral Surg 1976; 34: 19-23.
23. De Assis EA, Starck WJ, Epker B. Cephalometric analysis of profile nasal esthetics. Part III. Postoperativ changes after isolated superior repositioning. Int J Adult Orthod Orthognath 1996; 11: 279-288.
24. Ellis E; Hinton Rj. Histologic examination of the temporomandibular joint after mandibular advancement with and without rigid fixation: An experimental investigation in adult Maccaca mulatta. J Oral Maxillofac Syrg 1991; 49: 1316.
25. Ellis E. Condylar Positioning Devices for Orthodontic Surgery: Are They Necessary. J Oral Maxillofac Syrg 1994; 52: 536-552.
26. Epker BN, Fish LC. Surgical-orthodontic correction of open-bite deformity. Am J Orthod 1977; 71: 278-299.
27. Epker BN, Fish LC. Surgical superior repositioning of the maxilla: What to do with the mandible?. Am j Orthod 1980; 78: 164-191.
28. Epker B. Esthetic maxillofacial surgery. Lea und Febiger Verlag, Philadelphia, 1994.

29. Ermel, T., Hoffmann J.& Alfter G., et al.: Long-term stability of treatment results after upper jaw segmented osteotomy according to Schuchardt for correction of anterior open bite. *J Orofac Orthop* 1999, 60: 36-245
30. Freihofer HPM. Changes in nasal Profile after maxillary advancement in cleft and non-cleft patients. *J Maxillofac Surg* 1977; 5: 20-27.
31. Friede H, Kahnberg K-E, Adell R, Riedell A. Accuracy of cephalometric prediction in orthognathic surgery. *J Oral Maxillofac Surg* 1987; 45: 754-760.
32. Hogeman KE, Sarnäs KV, Willmar K. One-stage surgical and dental-orthopedic correction of bimaxillary and craniofacial dysostosis. In: *Fortschr Kiefer Gesichtschr*, Bd 18: 39-49, Thieme Verlag, Stuttgart, 1974.
33. Hui E, Hägg EUO, Tidman H. Soft tissue changes following maxillary osteotomies in cleft lip and palate and non-cleft Patients. *J Craniomaxillofac Surg* 1994; 22: 182-186.
34. Hullihen, S.R. Case of elongation of the under jaw and distortion of the face and neck, caused by a burn. *Dent.Cosmos*, Philadelphia. 42: 287-293, 1900. (Reprint des Artikels von 1849).
35. Jensen AC, Sinclair PM, Wolford LM. Soft tissue changes associated with double jaw surgery. *Am J Orthod Dentofac Orthop* 1992; 101: 266-275.
36. Kondo E. & Aoba, T.J. (2000). Nonsurgical and Nonextraction Treatment of Skeletal Class III Open Bite: Its Long-Term Stability. *American Journal of Orthodontics and Dentofacial Orthopedics*, 2000, 117: 267-287
37. Kondo E. & Arai, S.: Nonsurgical and Nonextraction Treatment of a Skeletal Class III Adult Patient with Severe Prognathic Mandible. *World Journal of Orthodontics*, 2005, .6: 233-247
38. Lanz, O.: Osteoplastische Resektion beider Oberkiefer nach Kocher. In Lücke, Rose. Vogel (Herg.): *Deutsche Zeitschrift für Chirurgie*. Leipzig, 1893
39. Lee DY, Bailey LJ, Proffit WR. Soft tissue changes after superior repositioning of the Maxilla with Le Fort I osteotomy: 5-years follow-up. *Int J Adult Orthod Orthognath* 1996; 11: 301-311.
40. Legan HL, Burstone GJ. Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 1980; 38: 744-51.
41. Lindorf HH. Funktionsstabile Tandem-Verschraubung der sagittalen Ramusosteotomie –Operationsthenik, neue Instrumente und Erfahrungen. *Dtsch Z Mund Kiefer Gesichts chir* 1984; 8: 367-373.
42. Lindorf HH. Sagittal ramus osteotomy with tandem screw fixation –technique and results. *J. Maxillofac Surg* 1986; 14: 311-316.
43. Liou, E.J.; Chen, P.H.; Wang, Y.C.; Yu, C.C.; Huang, C.S. & Chen, Y.R.: Surgery-First Accelerated Orthognathic Surgery: Orthodontic Guidelines and Setup For Model Surgery, *Journal of Oral and Maxillofacial Surgery*, 2011, 69: 771-780
44. O'Brien, K.; Wright, J.; Conboy, F.; Appelbe, P.; Bearn, D.; Caldwell, S.; Harrison, J.; Hussain, J.; Lewis, D.; Littlewood, S.; Mandall, N.; Morris, T.; Murray, A.; Oskouei, M.; Rudge, S.; Sandler, J.; Thiruvenkatachari, B.; Walsh, T. & Turbill, E.: Prospective, Multi-Center Study

of the Effectiveness of Orthodontic/Orthognathic Surgery Care in the United Kingdom, American Journal of Orthodontics and Dentofacial Orthopedics, 2009, 135: 709-714

45. Obwegeser H., Trauner R. Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien. Dtsch Zahn Mund Kieferheilk 1955; 23: 1-17.
46. Obwegeser H. The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. J Oral Surg 1957; 10: 687-692.
47. Obwegeser H. The indication for surgical correction of mandibular deformity by sagittal splitting technique. Br J Surg 1963; 1: 157-166.
48. Partsch C.: Ein neue Methode temporärer Gaumenresektion. Arch. Klein.Chir. 57: 874-858, 1898
49. Radney LJ, Jacobs JD. Soft tissue changes associated with surgical total maxillary intrusion. Am J Orthodont 1981; 80: 191-212.
50. Rekow ED, Speidel TM, Koenig RA. Location of the mandibular center of Autorotation in maxillary impaction surgery. Am J Orthod Dentofac Orthop 1993; 103: 530-536.
51. Rosen HM. Lip-nasal esthetics following Le Fort I Osteotomy. Plast Reconstr Surg 1988; 81: 171-182.
52. Schuchardt K. Formen des offenen Bisses und ihre operativen Behandlungsmöglichkeiten. Fortsch Kiefer Gesichtschir. Bd 1, Thieme Verlag, Stuttgart, 1955.
53. Stansbury CD, Evans CA, Miloro M, BeGole EA, Morris DE.: Stability of open bite correction with sagittal split osteotomy and closing rotation of the mandible. J Oral Maxillofac Surg 2010;68:149-159.
54. Togawa R, Iino S, Miyawaki S.: Skeletal Class III and open bite treated with bilateral sagittal split osteotomy and molar intrusion using titanium screws. Angle Orthod 2010, 80: 1176-184.
55. Watted N, Bill J. Untergesichtsverlängerung bei der Klasse-II-Dysgnathien: Neues von der kieferorthopädisch-kieferchirurgischen Therapie. Ästhet Zahnmed 1999; 4: 1992-300.
56. Watted N, Witt E, Bill J. Therapy Concept for the combined orthodontic-surgical Treatment of Angle Class II Deformities with Short-face Syndrome: New Aspects for surgical Lengthening of the Lower Face. Clin Orthod research, 2000; 3:78-93
57. Watted N, Bill J, Witt E, Reuther J. Lengthening of the lower face Angle class II patients with skeletal deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbourg, France 1999.
58. Westermark AH, Bystedt H, von Konow L, Sällström KO. Nasolabial morphology after Le Fort I Osteotomies. Int J Oral Maxillofac Surg 1991; 20: 25-30.
59. Williams EH, Evans DL, Barton WA, Williams BH. The effect of bite plane use on terminal hinge axis location, Angle Orthod. 1977; 47: 25-33.

60. Williamsone EH, Caves SA, Edenfield RJ, Morse PK. Cephalometric analysis: comparisons between maximum intercuspatation and centric relation, Am J Orthod 1978; 74: 672-677.
61. Wolford L.M., Epker B.N.: The combined anterior and posterior maxillary osteotomy a new technique. J. Oral Surg. 14: 123-127, 1975