

**Ausgabe:** face 1/2015

**Thema:** Tattoorentfernung mit zwei Lasern

**Autoren:** Dr. med. Johanna Koch, Dr. med. Melanie Schulz, Reinhard Gansel

---

## Literatur

1. Gopee, N., Warbritton, A., Miller, B., Couch, L., & Wamer, W. (2005). Response of mouse skin to tattooing: use of SKH-mice as a surrogate model for human tattooing. *Toxicol Appl Pharmacol* , 209.
2. Hogsberg, T., Jacobsen, N., Clausen, P., & Serup, J. (2013). Black tattoo inks induce reactive oxygen species production correlating with aggregation of pigment nanoparticles and product brand but not with the polycyclic aromatic hydrocarbon content. *Exp Dermatol* , 22 (7), 464-9.
3. IARC. (1996). Carbon black. *IARC Monogr Eval Carcinog Risks Hum* , 65, 149-262.
4. Ibrahimi, O., Sakamoto, F., Avram, M., & Anderson, R. (2011). Treatment of tattoo allergy with ablative fractional resurfacing: a novel paradigm for Tattoo removal. *J Am Acad Dermatol* , 64 (6), 1111-4.
5. Kim, J., Lee, J., Won, Y., Kim, J., & Lee, S. (2006). Titanium, a major constituent of blue ink, causes resistance to Nd-YAG (1064 nm) laser: results of animal experiments. *Acta Derm Venerol* , 86 (2), 110-3.
6. Landthaler, M. (3. April 2012). Prof. Dr. Tattoo und Krebsrisiko . (Süddeutsche.de, Interviewer)
7. Lucinda, T., Hazel, O., Joyce, L., & Hon, C. (2013). Successful Treatment of Tattoo-Induced Pseudolymphoma with Sequential Ablative Fractional Resurfacing Followed by Q-switched Nd:YAG 532 nm Laser. *J Cutan Aesthet Surg* , 6 (4), 226-8.
8. Marini, L., Kozarev, J., Grad, L., Jezersek, M., & Bencic, B. (2012). Fractional Er:YAG Skin Conditioning for Enhanced Efficacy of Nd:YAG Q-Switched Laser Tattoo Removal. *J Laser and Health Academy* , 1, 35-40.
9. Moehrle, M., Blaheta, H., & Ruck, P. (2001). Tattoo pigments mimics positive sentinel lymph node mimicking metastatic malignant melanoma. *Plast Reconstr Surg*, 203, 342-4.
10. Radmanesh, M., & Rafiei, Z. (2014). Combination of CO2 and Q-switched Nd:YAG lasers is more effective than Q-switched Nd:YAG laser alone for eyebrow tattoo removal. *J Cosmet Laser Ther* , 22, 1-4.

- 11. Regensburger, J., Lehner, K., Maisch, T., Vasold, R., Santarelli, F., Engel, E., et al. (2010). Tattoo inks contain polycyclic aromatic hydrocarbons that additionally generate deleterious singlet oxygen. *Exp Dermatol* , 19, e275-81.**
- 12. Reid, R., & Muller, S. (1980). Tattoo removal by CO laser dermabrasion. *Plast Reconstr Surg* , 65 (6), 717-28.**
- 13. Sanghavi, S., Dongre, A., & Khopkar, U. (2013). Tattoo reactions-An epidemic on surge:A report of 3 cases. *Indian J Dermatol* , 79 (2), 231-4.**
- 14. Seitz, A., Grunewald, S., Wagner, J., Simon, J., & Paasch, U. (2014). Fractional CO2 Laser is as effective as Q-switched ruby laser for the initial treatment of a traumatic tattoo. *J Cosmet Laser Ther* , 16 (6), 303-5.**
- 15. Valberg, P., Long, C., & Sax, S. (2006). Integrating studies on carcinogenic risk of carbon black: epidemiology, animal exposures, and mechanism of action. *J Occup Environ Med* , 48, 1291-1307.**
- 16. Worret, W., Raulin, C., Greve, B., & Hammes, S. (2009). Ausgelaufen, verblasst oder unerwünscht-So verschwinden Tätowierungen und Permanent-Make-up. *Ästhetische Dermatologie* , 5, 6-12.**