

Vorteile der Verwendung eines Bulkfill-Materials in der Kinderzahnheilkunde

Dr. Vicky Ehlers/Mainz

Dental Tribune Austrian Edition 4/2015

Literatur:

1. Fleming, G. J., Awan, M., Cooper, P. R., Sloan, A. J.: The potential of a resin-composite to be cured to a 4 mm depth. Dent Mater 24, 522 - 529 (2008)
2. Burgess, J., Cakir, D.: Comparative properties of low-shrinkage composite resins. Compend Contin Educ Dent 31, 10-15 (2010)
3. Roggendorf, M. J., Krämer, N., Appelt, A., Naumann, M., Frankenberger, R.: Marginal quality of flowable 4-mm base vs. conventionally layered resin composite. J Dent 39, 643 -647 (2011)
4. Ilie, N., Hickel, R.: Investigations on a methacrylate-based flowable composite based on the SDR™ technology. Dent Mater 27, 348 - 355 (2011)
5. Rullmann, I., Schattenberg, A., Marx, M., Willershausen, B., Ernst, C. P.: Photoelastic determination of polymerization shrinkage stress in low-shrinkage resin composites. Schweiz Monatsschr Zahnmed 122, 294 - 299 (2012)
6. Krämer, N., Frankenberger, R.: Compomers in restorative therapy of children: a literature review. Int J Paediatr Dent 17, 2 - 9 (2007)