

Literaturverzeichnis

Der dritte mesiale Wurzelkanal im zweiten Unterkiefermolaren

Preet Singh

Endodontie Journal 2/2015

- Ahmed, Abu-bakr, Yahia, & Ibrahim. (2007). Root and canal morphology of permanent mandibular molars in a Sudanese population. *Int Endod J*, 40(10), 766-771.
- Ballullaya, Vemuri, & Kumar. (2013). Variable permanent mandibular first molar: Review of literature. *J Conserv Dent*, 16(2), 99-110.
- Baugh, & Wallace. (2004). Middle mesial canal of the mandibular first molar: a case report and literature review. *J Endod*, 30(3), 185-186.
- Bergenholtz. (1974). Micro-organisms from necrotic pulp of traumatized teeth. *Odontol Revy*, 25(4), 347-358.
- Bystrom, & Sundqvist. (1981). Bacteriologic evaluation of the efficacy of mechanical root canal instrumentation in endodontic therapy. *Scand J Dent Res*, 89(4), 321-328.
- Friedman. (2002). Considerations and concepts of case selection in the management of post-treatment endodontic disease (treatment failure). *Endod Topics*, 1, 24.
- Hess. (1917). Zur Anatomie der Wurzelkanäle des menschlichen Gebisses mit Berücksichtigung der feinen Verzweigungen am Foramen apicale. *Schweiz Vjschr Zahnheilk*, 1, 38-43.
- Hess. (1925). *The Anatomy of the Root Canals of the Teeth of the permanent Dentition* (Vol. 1st edn.). London, UK: John Bale, Sons and Danielsson Ltd.
- Hülsmann, & Tulus. (2010). Der dritte mesiale Wurzelkanal in Uterkiefermolaren ("middle mesial"). *Endodontie*, 19, 123-131.
- Ingle, & Zeldow. (1958). An evaluation of mechanical instrumentation and the negative culture in endodontic therapy. *J Am Dent Assoc*, 57(4), 471-476.
- Kakehashi, Stanley, & Fitzgerald. (1965). The Effects of Surgical Exposures of Dental Pulp in Germ-Free and Conventional Laboratory Rats. *Oral Surg Oral Med Oral Pathol*, 20, 340-349.
- Manning. (1990a). Root canal anatomy of mandibular second molars. Part I. *Int Endod J*, 23(1), 34-39.
- Manning. (1990b). Root canal anatomy of mandibular second molars. Part II. C-shaped canals. *Int Endod J*, 23(1), 40-45.
- Moller, Fabricius, Dahlen, Ohman, & Heyden. (1981). Influence on periapical tissues of indigenous oral bacteria and necrotic pulp tissue in monkeys. *Scand J Dent Res*, 89(6), 475-484.
- Nosrat, Deschenes, Tordik, Hicks, & Fouad. (2015). Middle Mesial Canals in Mandibular Molars: Incidence and Related Factors. *J Endod*, 41(1), 28-32.
- Orstavik. (1996). Time-course and risk analyses of the development and healing of chronic apical periodontitis in man. *Int Endod J*, 29(3), 150-155.
- Peiris, Pitakotuwage, Takahashi, Sasaki, & Kanazawa. (2008). Root canal morphology of mandibular permanent molars at different ages. *Int Endod J*, 41(10), 828-835.
- Pomeranz, Eidelman, & Goldberg. (1981). Treatment considerations of the middle mesial canal of mandibular first and second molars. *J Endod*, 7(12), 565-568.
- Rollison, Barnett, & Stevens. (2002). Efficacy of bacterial removal from instrumented root canals in vitro related to instrumentation technique and size. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 94(3), 366-371.

- Siqueira, & Rocas. (2008). Clinical implications and microbiology of bacterial persistence after treatment procedures. *J Endod*, 34(11), 1291-1301 e1293.
- Tan, & Messer. (2002). The quality of apical canal preparation using hand and rotary instruments with specific criteria for enlargement based on initial apical file size. *J Endod*, 28(9), 658-664.
- Usman, Baumgartner, & Marshall. (2004). Influence of instrument size on root canal debridement. *J Endod*, 30(2), 110-112.
- Vertucci, & Williams. (1974). Root canal anatomy of the mandibular first molar. *J N J Dent Assoc*, 45(3), 27-28 passim.