

Ausgabe: face 3/2015

Thema: Früherkennung von Melanomen und automatisierte Analyse von Hautveränderungen mittels Elektrischer Impedanzspektroskopie

Autor: Dr. med. Emanuel von Kienlin

Literatur

1. Bay C, Kejs AM, Storm HH, Engholm G: Incidence and survival in patients with cutaneous melanoma by morphology, anatomical site and TNM stage: A Danish Population-based Register Study 1989-2011. *Cancer Epidemiol.* 2014 Nov 4; pii: S1877-7821(14)00188-X
2. Kardynal A, Olszewska M: Modern non-invasive diagnostic techniques in the detection of early cutaneous melanoma. *J Dermatol Case Rep.* 2014 Mar 31;8(1):1-8
3. Reinhold U: Moderne Diagnoseverfahren in der Dermato-Onkologie. *Spektrum der Dermatologie* 2013,2: 3-7
4. Birgersson U: Electrical Impedance of human skin and tissue alterations: Mathematical modeling and measurements. Thesis for doctoral degree. Karolinska Institutet 2012
5. Aberg P, Birgersson U, Elsner P, et al.: Electrical impedance spectroscopy and the diagnostic accuracy for malignant melanoma. *Exp. Dermatol* 2011; 20(8):648–652.
6. Mohr P, Birgersson U, Berking C, et al.: Electrical impedance spectroscopy as a potential adjunct diagnostic tool for cutaneous melanoma. *Skin Res Technol* 2013;19(2):75–83.
7. Malvey J, Hauschild A, Curiel-Lewandrowski C, et al.: Clinical performance of the Nevisense system in cutaneous melanoma detection: an international, multicentre, prospective and blinded clinical trial on efficacy and safety. *Br J Dermatol.* 2014 Nov;171(5):1099-107
8. SciBase AB (Accessed in 2014, at: <http://scibase.se/en/the-nevisense-product/the-eis-method/>)
9. Leiter U, Eigentler T, Garbe C: Epidemiology of skin cancer. *Adv Exp Med Biol.* 2014;810:120-40.