

## „Vorteile der KGIZ überwiegen bei Weitem“

**Beim AAO-Kongress referierte Prof. Dr. Roberto Justus zur „Entproteinisierung von Zahnschmelzoberflächen zur Verhinderung von White Spot Läsionen und Bracketklebeverlusten – eine Revolution im kieferorthopädischen Bonding“. KN traf den Direktor für Forschung der Abteilung KFO der Intercontinental University Mexico City zum Interview.**

### Literatur

1. Richter AE, Arruda AO, Peters MC, et al. Incidence of caries lesions among patients treated with comprehensive orthodontics. Am J Orthod Dentofacial Orthop. 2011;139:657–64.
2. Boersma JG, van der Veen MH, Lagerweij MD, et al. Caries prevalence measured with quantitative light-induced fluorescence after treatment with fixed orthodontic appliances: influencing factors. Caries Res. 2005;39:41–7.
3. Ogaard B. Prevalence of white spot lesions in 19-year olds: a study on untreated and orthodontically treated persons 5 years after treatment. Am J Orthod Dentofacial Orthop. 1989;96:423–7.
4. van der Veen MH, Mattousch T, Boersma JG. Longitudinal development of caries lesions after orthodontic treatment evaluated by quantitative light-induced fluorescence. Am J Orthod Dentofacial Orthop. 2007;131:223–8.
5. Marcusson A, Norevall LI, Persson M. White spot reduction when using a glass ionomer cement for bonding in orthodontics: a longitudinal and comparative study. Eur J Orthod. 1997;19:233–42.
6. Czochrowska EM, Ogaard B, Duschner H, et al. Cariostatic effect of a light-cured, resin-reinforced glass-ionomer for bonding orthodontic brackets *in vivo*. A combined study using microradiography and confocal laser scanning microscopy. J Orofac Orthop. 1998;59:265–73.
7. Wilson RM, Donly KJ. Demineralization around orthodontic brackets bonded with resin-modified glass ionomer cement and fluoride-releasing resin composite. J Pediatr Dent. 2001;23:255–9.
8. Sudjalim TR, Woods MG, Manton DJ, et al. Prevention of demineralization around orthodontic brackets *in vitro*. Am J Orthod Dentofacial Orthop. 2007;131:705.e1–9.
9. Chin MYH, Sandham A, Rumachik EN, et al. Fluoride release and cariostatic potential of orthodontic adhesives with and without daily fluoride rinsing. Am J Orthod Dentofacial Orthop. 2009;136:547–53.
10. Benson PE, Shah AA, Millett DT, et al. Fluorides, orthodontics and demineralization: a systematic review. J Orthod. 2005;32:102–14.
11. Rogers S, Chadwick B, Treasure E. Fluoride-containing orthodontic adhesives and decalcification in patients with fixed appliances: a systematic review. Am J Orthod Dentofacial Orthop. 2010;138:390.e1–8.
12. Silverstone LM, Saxton CA, Dogon IL, et al. Variation in the pattern of acid etching of human dental enamel examined by scanning electron microscopy. Caries Res. 1975;9:373–87.

13. Hobson RS, Rugg-Gunn AJ, Booth TA. Acid etch patterns on the buccal surface of human permanent teeth. *Arch Oral Biol.* 2002;47:407–12.
14. Hannig M, Bösmann K. Die Abrasivität des Pellicels unter klinischen Gesichtspunkten. *Dtsch Zahnärztl Z.* 1987;42:1015–20.
15. Cruz R, Rolla G. Deposition of alkali-soluble fluoride on enamel surface with or without pellicle. *Scand J Dent Res.* 1991;99:96–9.
16. Hannig M, Joiner A. The structure, function and properties of the acquired pellicle. In: Duckworth RM, editor. *The teeth and their environment, Monographs in oral science*, vol. 19. Basel: Karger; 2006. p. 29–64.
17. Reynolds IR. A review of direct orthodontic bonding. *Br J Orthod.* 1979;2:171–78.
18. Justus R, Cubero T, Ondarza R, et al. A new technique with sodium hypochlorite to increase bracket shear bond strength of fluoride-releasing resin-modified glass ionomer cements: comparing shear bond strength of two adhesive systems with enamel surface deproteinization before etching. *Semin Orthod.* 2010;16:66–75.
19. Larmour CJ, Stirrups DR. An ex vivo assessment of a resin-modified glass ionomer cement in relation to bonding technique. *J Orthod.* 2001;28:207–10.
20. Justus R. *Iatrogenic Effects of Orthodontic Treatment: Decision-making in Prevention, Diagnosis and Treatment.* Berlin: Springer International Publishing; 2015.
21. Hegarty DJ, Macfarlane TV. In vivo bracket retention comparison of a resin-modified bracket adhesive system after a year. *Am J Orthod Dentofacial Orthop.* 2002;121:496–501.
22. Bishara SE, VonWald L, Olsen ME, et al. Effect of time on the shear bond strength of glass ionomer and composite orthodontic adhesives. *Am J Orthod Dentofacial Orthop.* 1999;116:616–20.
23. Vivanco C. Effect of time on the shear bond strength of an ionomer and a composite: an ex-vivo study. MS Thesis, Intercontinental University, Mexico City; 2014.