

**Ausgabe:** KN 1/2-2016, S. 1ff.

**Thema:** Der progene Formenkreis – Therapie von Klasse III-Dysgnathien

**Autoren:** Dr. Heiko Goldbecher, Priv.-Doz. Dr. Dr. Robin Seeberger

---

## Literatur

- 1 Proffit WR, Fields HW, Jr., Moray LJ. Prevalence of malocclusion and orthodontic treatment need in the United States: estimates from the NHANES III survey. *The International journal of adult orthodontics and orthognathic surgery*. 1998;13(2):97-106.
- 2 Ramires T, Maia RA, Barone JR. Nasal cavity changes and the respiratory standard after maxillary expansion. *Brazilian journal of otorhinolaryngology*. 2008;74(5):763-9.
- 3 Hartgerink DV, Vig PS, Abbott DW. The effect of rapid maxillary expansion on nasal airway resistance. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*. 1987;92(5):381-9.
- 4 Seeberger R, Kater W, Schulte-Geers M, Thiele OC, Davids R, Hofele CH, et al. [Surgically assisted rapid maxillary expansion. Effects on the nasal airways and nasal septum]. *Hno*. 2010;58(8):806-11
- 5 Bell WH, Epker BN. Surgical-orthodontic expansion of the maxilla. *American journal of orthodontics*. 1976;70(5):517-28.
- 6 Wriedt S, Kunkel M, Zentner A, Wahlmann U.: Die chirurgisch unterstützte Gaumennahterweiterung, Reflexionsakustisch-volumetrische, morphometrische und sonographische Untersuchungen. *Fortschritte der Kieferorthopädie*. 2001; 62: 107-115.
- 7 Seeberger R, Kater W, Schulte-Geers M, Davids R, Freier K, Thiele O: Changes after surgically-assisted maxillary expansion (SARME) to the dentoalveolar, palatal and nasal structures by using tooth-borne distraction devices. *Br J Oral Maxillofac Surg*. 2011 Jul;49(5):381-5.
- 8 Tausche E, Deeb W, Hansen L, Hietschold V, Harzer W, Schneider M (2009): CT analysis of nasal volume changes after surgically-assisted rapid maxillary expansion. *J Orofac Orthop*;70(4):306-17
- 9 Timms DJ (1984): The reduction of nasal airway resistance by rapid maxillary expansion and its effect on respiratory disease. *J Laryngol Otol*; 98(4): 357-62
- 10 Seeberger R, Abe-Nickler D, Hoffmann J, Kunzmann K, Zingler S.: One-stage tooth-borne distraction versus two stage bone-borne distraction in surgically assisted maxillary expansion (SARME). *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2015 Dec;120(6):693-8.
- 11 Seeberger R, Gander E, Hoffmann J, Engel M.: Surgical management of cross-bites in orthognathic surgery: Surgically assisted rapid maxillary expansion (SARME) versus two-piece maxilla. *J Craniomaxillofac Surg*. 2015 Sep;43(7):1109-12.

- 12 Marchetti C, Pironi M, Bianchi A, Musci A: Surgically assisted rapid palatal expansion vs. segmental le Fort I osteotomy: transverse stability over a 2-year period. *J Craniomaxillofac Surg* 37:74-78, 2009.
- 13 Obwegeser, H., The Indications for Surgical Correction of Mandibular Deformity by the Sagittal Splitting Technique. *Br J Oral Surg*, 1964. 1: p. 157-71.
- 14 Hunsuck EE. A modified intraoral sagittal splitting technic for correction of mandibular prognathism. *J Oral Surg*. 1968;26:250–253.
- 15 Epker BN. Modifications in the sagittal osteotomy of the mandible. *J Oral Surg*. 1977;35:157–159.
- 16 Seeberger R, Asi Y, Thiele OC, Hoffmann J, Stucke K, Engel M.: Neurosensory alterations and function of the temporomandibular joint after high oblique sagittal split osteotomy: an alternative technique in orthognathic surgery. *Br J Oral Maxillofac Surg*. 2013 Sep;51(6):536-40.