

Ausgabe: Implantologie Jahrbuch 2016, S. 98-102

Thema: Bedingt herausnehmbare Versorgung des zahnlosen Oberkiefers auf sechs Implantaten

Autoren: Dr. med. dent. Kleanthis Manolakis, ZTM Diamantis Tsifoutakos, ZTM Georgios Nikolaou

Literatur

Bueno-Samper A, Hernandez-Aliaga M, Calvo-Guirado JL: The implant-supported milled bar overdenture: a literature review. *Med Oral Patol Oral Cir Bucal* (2010) 15:e375-378.

Heintze SD, Rousson V: Survival of zirconia- and metal-supported fixed dental prostheses: a systematic review. *Int J Prosthodont* (2010) 23:493-502.

Mericske-Stern R, Worni A: Optimal number of oral implants for fixed reconstructions: a review of the literature. *Eur J Oral Implantol* (2014) 7 Suppl 2:S133-153.

Mertens C, Steveling HG: Implant-supported fixed prostheses in the edentulous maxilla: 8-year prospective results. *Clin Oral Implants Res* (2011) 22:464-472.

Pomares C: A retrospective study of edentulous patients rehabilitated according to the 'all-on-four' or the 'all-on-six' immediate function concept using flapless computer-guided implant surgery. *Eur J Oral Implantol* (2010) 3:155-163.

Schimmel M, Srinivasan M, Herrmann FR, Muller F: Loading protocols for implant-supported overdentures in the edentulous jaw: a systematic review and meta-analysis. *Int J Oral Maxillofac Implants* (2014) 29 Suppl:271-286.

Deutsche Gesellschaft für Zahn-, Mund- und Kieferheilkunde: S3-Leitlinie: Implantatprothetische Versorgung des zahnlosen Oberkiefers (2013). Internet: <http://www.awmf.org/leitlinien/detail/II/083-010.html> [07.12.2014].