

Ausgabe: PN Parodontologie Nachrichten 3/16

Thema: **Vorsorge für die Mundgesundheit**

Autoren: Dr. Deborah Horch

Literatur

[1] Scheinin A, Mäkinen KK, Ylitalo K.: Turku sugar studies. V. Final report on the effect of sucrose, fructose and xylitol diets on the caries incidence in man, *Acta Odontol Scand.* 1976;34 (4):179-216

[2] Mäkinen, K.: Der steinige Weg von Xylitol bis zu seiner klinischen Anwendung, *J. Dentistry* 79(2000) pp.1352-1355

[3] Mäkinen, K.: Zuckeralkohole als Alternative zu Haushaltszucker mit besonderer Berücksichtigung von Xylitol, *Med Pract Prinz* 20(2011), pp. 302-320

[4] Mäkinen, K.: Xylitol-Kaugummis und Kariesrate, eine 40 monatige Kohortenstudie, *J Dent Res.* 74-12(1995), pp. 1904-1913

[5] Park, E. et al.: Xylitol, an anticaries agent, exhibits potent inhibition of inflammatory responses in human THP-1-derived macrophages infected with *Porphyromonas gingivalis*, *J Periodontol.* 2014 Jun, 85(6), pp. 212-223

[6] Uittamo J. et al.: Xylitol inhibits carcinogenic acetaldehyde production by *Candida* species, *Int J Cancer.* 2011 Oct 15;129(8), pp. 2038-2041

[7] The EFSA Journal (2008) 852, pp. 1-15: Xylitol chewing gum/pastilles and reduction of the risk of tooth decay, Scientific substantiation of a health claim related to xylitol chewing gum/pastilles and reduction the risk of tooth decay pursuant to Article 14 of Regulation (EC) No 1924/20061, Scientific Opinion of the Panel on Dietetic Products, Nutrition and Allergies (Question No EFSAQ- 2008-321) Adopted on 30 October 2008 by written procedure