

Ausgabe: KN 6-2016, S. 1ff.

Thema: Das PowerScope™ 2

Autoren: Dr. Andrew Hayes

Literatur

- Cacciatore, G., Alvetro, L., Defraia, E., Ghislanzoni, L. T., & Franchi, L. (2014). Active-treatment effects of the Forsus fatigue resistant device during comprehensive Class II correction in growing patients. *Korean J Orthod*, 44(3), 136-142.
doi:10.4041/kjod.2014.44.3.136
- Chen, J. Y., Will, L. A., & Niederman, R. (2002). Analysis of efficacy of functional appliances on mandibular growth. *Am J Orthod Dentofacial Orthop*, 122(5), 470-476.
doi:10.1067/mod.2002.126730
- Chhibber, A., Upadhyay, M., Uribe, F., & Nanda, R. (2013). Mechanism of Class II correction in prepubertal and postpubertal patients with Twin Force Bite Corrector. *Angle Orthod*, 83(4), 718-727. doi:10.2319/090412-709.1
- Dolce, C., McGorray, S. P., Brazeau, L., King, G. J., & Wheeler, T. T. (2007). Timing of Class II treatment: skeletal changes comparing 1-phase and 2-phase treatment. *Am J Orthod Dentofacial Orthop*, 132(4), 481-489. doi:10.1016/j.ajodo.2005.08.046
- Ehsani, S., Nebbe, B., Normando, D., Lagravere, M. O., & Flores-Mir, C. (2015). Short-term treatment effects produced by the Twin-block appliance: a systematic review and meta-analysis. *Eur J Orthod*, 37(2), 170-176. doi:10.1093/ejo/cju030
- Fontana, M., Cozzani, M., & Caprioglio, A. (2012). Soft tissue, skeletal and dentoalveolar changes following conventional anchorage molar distalization therapy in class II non-growing subjects: a multicentric retrospective study. *Prog Orthod*, 13(1), 30-41.
doi:10.1016/j.pio.2011.07.002
- Harrison, J. E., O'Brien, K. D., & Worthington, H. V. (2007). Orthodontic treatment for prominent upper front teeth in children. *Cochrane Database Syst Rev*(3), CD003452.
doi:10.1002/14651858.CD003452.pub2
- Jones, G., Buschang, P. H., Kim, K. B., & Oliver, D. R. (2008). Class II non-extraction patients treated with the Forsus Fatigue Resistant Device versus intermaxillary elastics. *Angle Orthod*, 78(2), 332-338. doi:10.2319/030607-115.1
- Manni, A., Pasini, M., & Mauro, C. (2012). Comparison between Herbst appliances with or without miniscrew anchorage. *Dent Res J (Isfahan)*, 9(Suppl 2), S216-221.
doi:10.4103/1735-3327.109762
- O'Brien, K., Wright, J., Conboy, F., Sanjie, Y., Mandall, N., Chadwick, S., . . . Shaw, I. (2003). Effectiveness of early orthodontic treatment with the Twin-block appliance: a multicenter, randomized, controlled trial. Part 1: Dental and skeletal effects. *Am J Orthod Dentofacial Orthop*, 124(3), 234-243; quiz 339.
doi:10.1016/S0889540603003524
- Tulloch, J. F., Proffit, W. R., & Phillips, C. (2004). Outcomes in a 2-phase randomized clinical trial of early Class II treatment. *Am J Orthod Dentofacial Orthop*, 125(6), 657-667. doi:10.1016/S088954060400160X
- Vaid, N., Doshi, V., & Vandekar, M. (2014). Class II treatment with functional appliances: A meta-analysis of short-term treatment effects. *Seminars in Orthodontics*, 324-338.

Weschler, D., & Pancherz, H. (2005). Efficiency of three mandibular anchorage forms in Herbst treatment: a cephalometric investigation. *Angle Orthod*, 75(1), 23-27.
doi:10.1043/0003-3219(2005)075<0023:EOTMAF>2.0.CO;2