

Ausgabe: Dental Tribune German Edition 2/2017, S. 8-9

Thema: Behandlung eines zentralen avitalen Schneidezahns mit offenem Apex mittels Anwendung von MTA-basiertem Reparaturmaterial

Autoren: Mario Luis Zuolo, DDS, MSc. und Arthur de Siqueira Zuolo, DDS, MSc, São Paulo, Brasilien

Literatur

- (1) Trope M. Treatment of immature teeth with non-vital pulps and apical periodontitis. *EndodTopics*.2006;14:51-59.
- (2) Rafter M.Apexification: a review.*DentTraumatol*.2005;21(1):1-8.
- (3) Granath LE. Some notes on the treatment of traumatized incisors in children. *OdontRev*.1959;10:272.
- (4) Morfis AS, Siskos G. Apexification with the use of calcium hydroxide: a clinical study.*JClinPediatrDent*.1991;16(1):13-19.
- (5) KleierDJ, Barr ES. A study of endodontically apexified teeth.*EndodDentTraumatol*. 1991;7(3):112-117.
- (6) WaliaT, Chawla HS, Gauba K. Management of wide open apices in non-vital permanent teeth with Ca(OH)₂ paste. *J ClinPediatrDent*. 2000;25(1):51-56.
- (7) Dominguez Reyes A, Munoz Munoz L, Aznar Martin T. Study of calcium hydroxide apexification in 26 young permanent incisors. *DentTraumatol*.2005;21(3):141-145.
- (8) Sübay RK, Kayataş M. Dens invaginatus in an immature maxillary lateral incisor: a case report of complex endodontic treatment. *Oral Surg Oral Med Oral Pathol Oral RadiolEndod*.2006;102(2):e37-41.
- (9) El-Meligy OA, Avery DR. Comparison of apexification with mineral trioxide aggregate and calcium hydroxide. *PediatrDent*. 2006;28(3):248-253.
- (10) Andreasen JO, Farik B, Munksgaard EC. Long-term calcium hydroxide as a root Kanal dressing may increase risk of root fracture.*DentTraumatol*.2002;18(3):134-137.
- (11) Andreasen JO, Munksgaard EC, Bakland LK. Comparison of fracture resistance in root Kanals of immature sheep teeth after filling with calcium hydroxide or MTA. *DentTraumatol*. 2006;22(3):154-156.
- (12) Heling I, Lustmann J, Hover R, Bichacho N. Complications of apexification resulting from poor patient compliance: report of case. *ASDC J DentChild*.1999;66(6):415-418.
- (13) Torabinejad M, Chivian N. Clinical applications of mineral trioxide aggregate. *J Endod*.1999;25(3):197-205.
- (14) Torabinejad M, Hong CU, McDonald F, Pitt Ford TR. Physical and chemical properties of a new root-end filling Material. *J Endod*. 1995;21(7):349-353.
- (15) Shabahang S, Torabinejad M, Boyne PP, Abedi H, McMillan P. A comparative study of root-end induction using osteogenic protein-1, calcium hydroxide, and mineral trioxide aggregate in dogs. *J Endod*.1999;25(1):1-5.

- (16) Simon S, Rilliard F, Berdal A, Machtou P. The use of mineral trioxide aggregate in one-visit apexification treatment: a prospective study. *IntEndod J.*2007;40(3):186–197.
- (17) Steinig TH, Regan JD, Gutmann JL. The use and predictable placement of mineral trioxide aggregate in one-visit apexification cases. *AustEndod J.*2003;29(1):34–42.
- (18) Witherspoon DE, Small JC, Regan JD, Nunn M. Retrospective analysis of open apex teeth obturated with mineral trioxide aggregate. *J Endod.*2008;34(10):1171-1176.
- (19) Mente J, Hage N, Pfefferle T, Koch MJ, Dreyhaupt J, Staehle HJ, Friedman S. Mineral trioxide aggregate apical plugs in teeth with open apical foramina: a retrospective analysis of treatment outcome. *J Endod.*2009;35(10):1354–1358.
- (20) Nayar S, Bishop K, Alani A. A report on the clinical and radio-graphic outcomes of 38 cases of apexification with mineral trioxide aggregate. *EurJProsthodontRestorDent.* 2009; 17(4): 150–156.
- (21) Holden DT, Schwartz SA, Kirkpatrick TC, Schindler WG. Clinical outcomes of artificial root-end barriers with mineral trioxide aggregate in teeth with immature apices. *JEndod.*2008; 34(7):812–817.
- (22) Seltzer S, Sinai I, August D. Periodontal effects of root perfora- tions before and during endodontic procedures. *J Dent Res.* 1970;49(2):332–339.
- (23) deChevigny, Dao TT, Basrani BR, Marquis V, Farzaneh M, Abitbol S, Friedman S. Treatment outcome in endodontics: the Toronto study — phases 3 and 4: orthograde retreatment. *J Endod.* 2008;34(2):131-137.
- (24) Angelus. MTA REPAIR HP. <http://angelus.ind.br/MTA-REPAIR-HP-292.html>. Accessed April 4, 2016.
- (25) Molven O, Halse A, Grung B. Incomplete healing (scar tissue) after periapical surgery — radiographic findings 8 to 12 years after treatment. *J Endod.*1996;2(5):264-268.