

Minimalinvasiv implantieren – einfach und praktikabel

Dr. Lara Müller, Dr. Frank Kistler, Dr. Georg Bayer/Landsberg am Lech

Implantologie Journal 1/2011

Literatur:

- [1] Pierrisnard L, Renouard F, Renault P, Barquins M., Influence of implant length and bicortical anchorage on implant stress distribution. Clin Implant Dent Relat Res. 2003;5(4):254-62
- [2] Baelum V, Ellegaard B. Implant survival in periodontally compromised patients. J Periodontol. 2004 Oct;75(10):1404-12.
- [3] Rangert B Mechanical and biomechanical guidelines for the use of Brånemark System--clinical studies.. Aust Prosthodont J. 1993;7 Suppl:45-9. Review.
- [4] Adell R, Eriksson B, Lekholm U, Brånemark PI, Jemt T. Long-term follow-up study of osseointegrated implants in the treatment of totally edentulous jaws. Int J Oral Maxillofac Implants. 1990 Winter;5(4):347-59.
- [5] Bahat O. Brånemark system implants in the posterior maxilla: clinical study of 660 implants followed for 5 to 12 years. Int J Oral Maxillofac Implants. 2000 Sep-Oct;15(5):646-53.
- [6] Becker W, Becker BE, Alsuwyed A, Al-Mubarak S. Long-term evaluation of 282 implants in maxillary and mandibular molar positions: a prospective study. J Periodontol. 1999 Aug;70(8):896-901.
- [7] Davarpanah M, Martinez H, Celletti R, Alcoforado G, Tecucianu JF, Etienne D Osseotite implant: 3-year prospective multicenter evaluation. Clin Implant Dent Relat Res. 2001;3(2):111-8.
- [8] Pierrisnard L, Renouard F, Renault P, Barquins M Influence of implant length and bicortical anchorage on implant stress distribution. Clin Implant Dent Relat Res. 2003;5(4):254-62.
- [9] Bernard JP, Szmukler-Moncler S, Pessotto S, Vazquez L, Belser UC. The anchorage of Brånemark and ITI implants of different lengths. I. An experimental study in the canine mandible. Clin. Oral Implants Res. 2003 ct.14 (5):593-600
- [10] Tawil G, Younan R. Clinical evaluation of short, machined-surface implants followed for 12 to 92 months. Int J Oral Maxillofac Implants. 2003 Nov-Dec;18(6):894-901.
- [11] Renouard F, Nisand D Short implants in the severely resorbed maxilla: a 2-year retrospective clinical study. Clin Implant Dent Relat Res. 2005;7 Suppl 1:S104-10.
- [12] Stellingsma K, Raghoobar GM, Meijer HJ, Stegenga B. The extremely resorbed mandible: a comparative prospective study of 2-year results with 3 treatment strategies. Int J Oral Maxillofac Implants. 2004 Jul-Aug;19(4):563-77.
- [13] Prof. Dr. Mauro Marincola/Rom, Vincent Morgan, DMD/Boston, Prof. Andrea Cicconetti/Rom, Jessica Betz/Bingen Einsatz von Kurzimplantaten bei Hypästhesie des N. alveolaris inferior Oral. Journal 3/2009 Aug; 24-30