

Indikationsabhängige Material- und Behandlungstechnikauswahl für den ästhetischen Frontzahnersatz

Autoren: Dr. Steffen Kistler, Marion Bayer, Dr. Frank Kistler, Priv.-Doz. Dr. Jörg Neugebauer

Literatur:

1. Buser D, Martin W, Belser UC. Optimizing esthetics for implant restorations in the anterior maxilla: anatomic and surgical considerations. *Int J Oral Maxillofac Implants* 2004; 19 Suppl: 43-61.
2. Buser D, Chen ST, Weber HP, Belser UC. Early implant placement following single-tooth extraction in the esthetic zone: biologic rationale and surgical procedures. *Int J Periodontics Restorative Dent* 2008; 28: 441-451.
3. Darby I, Chen ST, Buser D. Ridge preservation techniques for implant therapy. *The International journal of oral & maxillofacial implants* 2009; 24 Suppl: 260-271.
4. De Rouck T, Eghbali R, Collys K, De Bruyn H, Cosyn J. The gingival biotype revisited: transparency of the periodontal probe through the gingival margin as a method to discriminate thin from thick gingiva. *Journal of Clinical Periodontology* 2009; 36: 428-433.
5. Degidi M, Artese L, Scarano A, Perrotti V, Gehrke P, Piattelli A. Inflammatory infiltrate, microvessel density, nitric oxide synthase expression, vascular endothelial growth factor expression, and proliferative activity in peri-implant soft tissues around titanium and zirconium oxide healing caps. *J Periodontol* 2006; 77: 73-80.
6. Degidi M, Iezzi G, Scarano A, Piattelli A. Immediately loaded titanium implant with a tissue-stabilizing/maintaining design ('beyond platform switch') retrieved from man after 4 weeks: a histological and histomorphometrical evaluation. A case report. *Clin Oral Implants Res* 2008; 19: 276-282.
7. Fickl S, Zuh O, Wachtel H, Kebschull M, Hurzeler MB. Hard tissue alterations after socket preservation with additional buccal overbuilding: a study in the beagle dog. *J Clin Periodontol* 2009; 36: 898-904.
8. Furhauser R, Florescu D, Benesch T, Haas R, Mailath G, Watzek G. Evaluation of soft tissue around single-tooth implant crowns: the pink esthetic score. *Clin Oral Implants Res* 2005; 16: 639-644.
9. Garg AK. The Atlantis Components Abutment: simplifying the tooth implant procedure. *Dental implantology update* 2002; 13: 65-70.
10. Haessler D, Kornmann F. Implantatfreilegung mit gleichzeitiger Papillenrekonstruktion. *Quintessenz* 1998; 4: 763-768.
11. Kerstein RB, Radke J. A comparison of fabrication precision and mechanical reliability of 2 zirconia implant abutments. *The International journal of oral & maxillofacial implants* 2008; 23: 1029-1036.
12. Kuhlke KL, Drennon DG. An alternative to the anterior single-tooth removable partial denture. *J Int Assoc Dent Child* 1977; 8: 11-15.
13. Mijiritsky E, Mardinger O, Mazor Z, Chaushu G. Immediate provisionalization of single-tooth implants in fresh-extraction sites at the maxillary esthetic zone: up to 6 years of follow-up. *Implant Dent* 2009; 18: 326-333.
14. Neugebauer J, Jozsa M, Kubler A. [Antimicrobial photodynamic therapy for prevention of alveolar osteitis and post-extraction pain]. *Mund Kiefer Gesichtschir* 2004; 8: 350-355.
15. Nkenke E, Radespiel-Troger M, Wiltfang J, Schultze-Mosgau S, Winkler G, Neukam FW. Morbidity of harvesting of retromolar bone grafts: a prospective study. *Clinical oral implants research* 2002; 13: 514-521.
16. Rothamel D, Schwarz F, Herten M, Chiriac G, Pakravan N, Sager M, Becker J. [Dimensional ridge alterations following tooth extraction. An experimental study in the dog]. *Mund-, Kiefer- und Gesichtschirurgie : MKG* 2007; 11: 89-97.
17. Rothamel D, Schwarz F, Herten M, Engelhardt E, Donath K, Kuehn P, Becker J. Dimensional ridge alterations following socket preservation using a nanocrystalline hydroxyapatite paste: a histomorphometrical study in dogs. *Int J Oral Maxillofac Surg* 2008; 37: 741-747.
18. Salama H, Salama M, Garber D, Adar P. Developing optimal peri-implant papillae within the esthetic zone: guided soft tissue augmentation. *J Esthet Dent* 1995; 7: 125-129.
19. Schlee M, Esposito M. Aesthetic and patient preference using a bone substitute to preserve extraction sockets under pontics. A cross-sectional survey. *European journal of oral implantology* 2009; 2: 209-217.
20. Schmidlin PR, Jung RE, Schug J. [Prevention of alveolar ridge resorption after tooth extraction--a review]. *Schweiz Monatsschr Zahnmed* 2004; 114: 328-336.
21. Schulte W, Kleineikenscheidt H, Lindner K, Schareyka R. [The Tubingen immediate implant in clinical studies]. *Deutsche zahnärztliche Zeitschrift* 1978; 33: 348-359.
22. Stimmelmayr M, Allen EP, Reichert TE, Iglhaut G. Use of a combination epithelialized-subepithelial connective tissue graft for closure and soft tissue augmentation of an extraction site following ridge preservation or implant placement: description of a technique. *The International journal of periodontics & restorative dentistry* 2010; 30: 375-381.
23. Tarnow DP, Magner AW, Fletcher P. The effect of the distance from the contact point to the crest of bone on the presence or absence of the interproximal dental papilla. *J Periodontol* 1992; 63: 995-996.
24. Ten Heggeler JM, Slot DE, Van der Weijden GA. Effect of socket preservation therapies following tooth extraction in non-molar regions in humans: a systematic review. *Clinical oral implants research* 2010.
25. Tischler M, Misch CE. Extraction site bone grafting in general dentistry. Review of applications and principles. *Dentistry today* 2004; 23: 108-113.

26. Weng D, Nagata MJ, Bell M, Bosco AF, de Melo LG, Richter EJ. Influence of microgap location and configuration on the periimplant bone morphology in submerged implants. An experimental study in dogs. *Clin Oral Implants Res* 2008; 19: 1141-1147.
27. Wheeler SL. Implant complications in the esthetic zone. *J Oral Maxillofac Surg* 2007; 65: 93-102.