

Ausgabe: face 2/2016

Thema: Lippenaugmentation mit der neuen Fillergeneration Agarose-Gel

Autor: Dr. med. Arna Shab

Literatur

1. Scarano A. Ringiovanimento dei tessuti molli periorali con agarose gel. *Dent Clin* 2009;2:5-13.
2. Scarano A, Carinci F, Piattelli A. Lip augmentation with a new filler (agarose gel): a 3-year follow-up study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009 Aug;108(2):e11-5.
3. Cho YS, Hong ST, Choi KH, Chang YH, Chung AS. Chemo-preventive activity of porphyrin derivatives against 6-sulfooxymethylbenzo[a]pyrene mutagenicity. *Asian Pac J Cancer Prev* 2000;1:311-7.
4. Marczylo T, Arimoto-Kobayashi S, Hayatsu H. Protection against Trp-P-2 mutagenicity by purpurin: mechanism of in vitro antimutagenesis. *Mutagenesis* 2000;15:223-8.
5. Naziruddin B, Durriya S, Phelan D, Duffy BF, Olack B, Smith D, et al. HLA antibodies present in the sera of sensitized patients awaiting renal transplant are also reactive to swine leukocyte antigens. *Transplantation* 1998;66:1074-80.
6. Gu Y, Tabata Y, Kawakami Y, Balamurugan AN, Hori H, Nagata N, et al. Development of a new method to induce angiogenesis at subcutaneous site of streptozotocin-induced diabetic rats for islet transplantation. *Cell Transplant* 2001;10: 453-7.
7. Duffy DM. Complications of fillers: overview. *Dermatol Surg* 2005;31:1626-33.
8. Cohen JL. Understanding, avoiding, and managing dermal filler complications. *Dermatol Surg* 2008;34:S92-9.
9. Fernández-Cossío S, León-Mateos A, Sampedro FG, Oreja MT. Biocompatibility of agarose gel as a dermal filler: histologic evaluation of subcutaneous implants. *Plast Reconstr Surg.* 2007 Oct;120(5):1161-9.
10. Tabata M, Shimoda T, Sugihara K, Ogomi D, Serizawa T, Akashi M. Osteoconductive and hemostatic properties of apatite formed on/in agarose gel as a bone-grafting material. *J Biomed Mater Res B Appl Biomater.* 2003 Nov 15;67(2):680-8.